


## 5220 Arborescent matorral with Zyziphus

### 1. National level

Biogeographical regions and/or marine regions concerned within the Member State: **MED**



map-distribution

### 2. Biogeographical or marine level

2.1 Biogeographical region or marine region: **MEDITERRANEAN**

#### 2.2 Published sources and/or websites:

Bartolomé, C., J. Álvarez, J. Vaquero, M. Costa, M.A. Casermeiro, J. Giraldo & J. Zamora (2005). Los tipos de hábitat de interés comunitario de España. Guía básica. Dirección General para la Biodiversidad, Ministerio de Medio Ambiente.

Ministerio de Medio Ambiente. (2003). Atlas y manual de los hábitat de España. Dirección General de Conservación de la Naturaleza, Ministerio de Medio Ambiente.

#### 2.3 Range of the habitat type in the biogeographical region or marine region

- |  |   |
|--|---|
| 2.3.1 Surface area of range in km2:            | 5567  |
| 2.3.2 Date of range determination:             | 1997-2006   |
| 2.3.3 Quality of data concerning range:        | Moderate e.g. based on partial data with some extrapolation |
| 2.3.4 Range trend:                             | Unknown (X)   |
| 2.3.5 Range trend magnitude in km2 (optional): |   |
| 2.3.6 Range trend period:                      |   |
| 2.3.7 Reasons for reported trend:              |   |
| and/or specify                                 |   |

#### 2.4 Area covered by habitat type in the biogeographical region or marine region

- |   |  |
|---|--|
| 2.4.1 Surface area of the habitat type (km2): | 79   |
| 2.4.2 Date of area estimation:                | 1997-2006  |
| 2.4.3 Method used for area estimation:        | Ground based survey (based on field mapping, possibly using stratified random sa |
| 2.4.4 Quality of data on area:                | Good e.g based on extensive surveys  |
| 2.4.5 Area trend:                             | Unknown (X)  |
| 2.4.6 Area trend magnitude (km2):             | 0  |

5220 Arborescent matorral with Zyziphus

2.4.7 Area trend period:

2.4.8 Reasons for reported trend:

and/or specify:

2.4.9 Justification of % thresholds for trends (optional):

2.4.10 Main pressures: 100 - Cultivation  
170 - Animal breeding

2.4.11 Threats 100 - Cultivation  
170 - Animal breeding

2.5 Complementary information

2.5.1 Favourable reference range (km2): 0

2.5.2 Favourable reference area (km2): 0

2.5.3 Typical Species: *Asparagus albus*, *Asparagus stipularis*, *Calicotome infesta* ssp. *intermedia*, *Chamaerops humilis*, *Launaea arborescens*, *Lycium intricatum*, *Maytenus senegalensis* ssp. *Europaea*, *Periploca angustifolia*, *Periploca angustifolia* ssp. *laevigata*  
*Withania frutescens*, *Rhamnus oleoides* ssp. *angustifolia*, *Withania frutescens*, *Ziziphus lotus*

2.5.4 Typical species assessment:

2.5.5 Other relevant information (optional):

Conclusion	Biogeographical or marine level	Conclusions within Natura 2000 sites (optional)
Conclusions: (2.3) Range:	Unknown (XX)	
Conclusions: (2.4) Area:	Unknown (XX)	
Conclusions: (2.5) Structure and function, including typical species:	Unknown (XX)	
Conclusions: Future prospects:	Unknown (XX)	
Conclusions: Overall assessment:	Unknown (XX)	