

Helianthemum bystropogophyllum

1. National level

Biogeographical regions and/or marine regions concerned within the Member State: **MAC**

map-range

map-distribution



map-favourable-range



2. Biogeographical or marine level

2.1 Biogeographical region or marine region: **MACARONESIAN**

2.2 Published sources and/or websites:

Beltrán E., Wildpret W., León C., García A. & A. Reyes (1999). Libro Rojo de la Flora Canaria contenida en la Directiva-Hábitats Europea. Ministerio de Medio Ambiente. 694 pp.

Bañares A., Blanca G., Güemes J., Moreno J.C. & Ortiz S., eds. (2003). Atlas y Libro Rojo de la Flora Vascular Amenazada de España. Dirección General de Conservación de la Naturaleza. Madrid. 1072 pp.

http://www.mma.es/porta/cciones/biodiversidad/inventarios/inb/flora_vascular/pdf/223.pdf

2.3 Range of the species type in the biogeographic region or marine region

2.3.1 Surface area of species range in km2: 100

2.3.2 Date of range determination: 2003

2.3.3 Quality of data concerning range:

2.3.4 Range trend: Unknown (X)

Helianthemum bystropogophyllum

2.3.5 Range trend magnitude in km2 (optional):	
2.3.6 Range trend period:	
2.3.7 Reasons for reported trend:	Unknown
and/or specify	
2.4 Population of the species in the biogeographic region or marine region	
2.4.1 Population size estimation:	
Population size estimation (minimum)	Population size estimation (maximum)
32	Population units
	Number of individuals
2.4.2 Date of population estimation:	2003
2.4.3 Methods used for population estimation:	From comprehensive inventory
2.4.4 Quality of data on area:	Good e.g based on extensive surveys
2.4.5 Population trend:	Unknown (X)
2.4.6 Population trend magnitude (km2):	
2.4.7 Population trend period:	
2.4.8 Reasons for reported trend:	Unknown
and/or specify:	
2.4.9 Justification of % thresholds for trends (optional):	
2.4.10 Main pressures:	965 - predation
2.4.11 Threats	965 - predation
2,5 Habitat for the species in the biogeographic region or marine region	
2.5.1 Habitats for the species:	Aparece principalmente en taliscas y poyatas, en sustratos de ignimbritas fonolíti
2.5.2 Area estimation (km2):	100
2.5.3 Date of estimation:	2007
2.5.4 Quality of the data:	Moderate e.g. based on partial data with some extrapolation
2.5.5 Trend of the habitat:	Unknown (X)
2.5.6 Trend period:	1970-2006
2.5.7 Reasons for reported trend:	Unknown
Other (specify):	
2.6 Future prospects for the species:	Bad prospects - species likely to be become extinct in the biogeographical region
2.7 Complementary information	
2.7.1 Favourable reference range (km2):	Less than
2.7.2 Favourable reference population:	More than
2.7.3 Suitable habitat for the species (km2):	25
2.7.4 Other relevant information (optional):	La especie no tiene problemas para reproducirse, florece y produce semillas en c
Conclusion	Biogeographical or marine level
Conclusions: (2.3) Range:	Bad (U2)
Conclusions: (2.4) Population:	Bad (U2)
Conclusions: (2.5) Habitat for the species:	Inadequate and deteriorating (U1-)
Conclusions: (2.6) Future prospects:	Bad and deteriorating (U2-)
Conclusions: Overall assessment:	Bad (U2)
	Conclusions within Natura 2000 sites (optional)