

Report on the main results of the surveillance under article 11 for annex II, IV and V species (Annex B)

Sambucus palmensis

1. National level

Biogeographical regions and/or marine regions concerned within the Member State: **MAC**

map-range

map-distribution



map-favourable-range



2. Biogeographical or marine level

2.1 Biogeographical region or marine region: **MACARONESIAN**

2.2 Published sources and/or websites:

Beltrán E., Wildpret W., León C., García A. & A. Reyes (1999). Libro Rojo de la Flora Canaria contenida en la Directiva-Hábitats Europea. Ministerio de Medio Ambiente. 694 pp.

Bañares A., Blanca G., Güemes J., Moreno J.C. & Ortiz S., eds. (2003). Atlas y Libro Rojo de la Flora Vascular Amenazada de España. Dirección General de Conservación de la Naturaleza. Madrid. 1072 pp

http://www.mma.es/portal/secciones/biodiversidad/inventarios/inb/flora_vascular/pdf/141.pdf

2.3 Range of the species type in the biogeographic region or marine region

2.3.1 Surface area of species range in km2: 1400

2.3.2 Date of range determination: 2003

2.3.3 Quality of data concerning range:

2.3.4 Range trend: Increasing (+)

Sambucus palmensis

- 2.3.5 Range trend magnitude in km2 (optional):
- 2.3.6 Range trend period: 1984-2003
- 2.3.7 Reasons for reported trend: Direct human influence (restoration, deterioration, destruction)
and/or specify

2.4 Population of the species in the biogeographic region or marine region

- 2.4.1 Population size estimation:
- Population size estimation (minimum)

Population size estimation (maximum)

Population units

340

340

Number of individuals
- 2.4.2 Date of population estimation: 2003
- 2.4.3 Methods used for population estimation: From comprehensive inventory
- 2.4.4 Quality of data on area: Good e.g based on extensive surveys
- 2.4.5 Population trend: Increasing (+)
- 2.4.6 Population trend magnitude (km2):
- 2.4.7 Population trend period: 1985-2003
- 2.4.8 Reasons for reported trend: Other (specify)
and/or specify:
- 2.4.9 Justification of % thresholds for trends (optional):
- 2.4.10 Main pressures: 850 Modification of hydrographic functioning, general
965 - predation
- 2.4.11 Threats 965 - predation

2,5 Habitat for the species in the biogeographic region or marine region

- 2.5.1 Habitats for the species: Se desarrolla entre los 600-100 m.s.m. Sus mejores manifestaciones suelen prese
- 2.5.2 Area estimation (km2): 825
- 2.5.3 Date of estimation: 2007
- 2.5.4 Quality of the data: Moderate e.g. based on partial data with some extrapolation
- 2.5.5 Trend of the habitat: Stable (=)
- 2.5.6 Trend period: 1970-2006
- 2.5.7 Reasons for reported trend: NaturalProcesses
Other (specify):

- 2.6 Future prospects for the species: Good prospects - species expected to survive and prosper

2.7 Complementary information

- 2.7.1 Favourable reference range (km2): 275
- 2.7.2 Favourable reference population:
- 2.7.3 Suitable habitat for the species (km2): 275
- 2.7.4 Other relevant information (optional): Favourable reference area: 100 individuals/ population Las poblaciones naturales

Conclusion	Biogeographical or marine level	Conclusions within Natura 2000 sites (optional)
Conclusions: (2.3) Range:	Inadequate but improving (U1+)	
Conclusions: (2.4) Population:	Inadequate but improving (U1+)	
Conclusions: (2.5) Habitat for the species:	Inadequate but improving (U1+)	
Conclusions: (2.6) Future prospects:	Inadequate but improving (U1+)	

Sambucus palmensis

Conclusions: Overall assessment:

Inadequate but improving (U1+)