

Sideritis discolor

1. National level

Biogeographical regions and/or marine regions concerned within the Member State: **MAC**

map-range

map-distribution



map-favourable-range



2. Biogeographical or marine level

2.1 Biogeographical region or marine region: **MACARONESIAN**

2.2 Published sources and/or websites:

Beltrán E., Wildpret W., León C., García A. & A. Reyes, 1999. Libro Rojo de la Flora Canaria contenida en la Directiva-Hábitats Europea. Ministerio de Medio Ambiente. 694 pp.

Bañares A., Blanca G., Güemes J., Moreno J.C. & Ortiz S., eds. (2003). Atlas y Libro Rojo de la Flora Vascular Amenazada de España. Dirección General de Conservación de la Naturaleza. Madrid. 1072 pp

http://www.mma.es/porta1/secciones/biodiversidad/inventarios/inb/flora_vascular/pdf/737.pdf

2.3 Range of the species type in the biogeographic region or marine region

2.3.1 Surface area of species range in km2: 100

2.3.2 Date of range determination: 2003

2.3.3 Quality of data concerning range:

2.3.4 Range trend: Stable (=)

Sideritis discolor

2.3.5	Range trend magnitude in km2 (optional):	
2.3.6	Range trend period:	1992-2003
2.3.7	Reasons for reported trend:	Improved knowledge/more accurate data
	and/or specify	
2.4	Population of the species in the biogeographic region or marine region	
2.4.1	Population size estimation:	
	Population size estimation (minimum)	Population size estimation (maximum)
	104	104
		Population units
		Number of individuals
2.4.2	Date of population estimation:	2003
2.4.3	Methods used for population estimation:	From comprehensive inventory
2.4.4	Quality of data on area:	Good e.g based on extensive surveys
2.4.5	Population trend:	Stable (=)
2.4.6	Population trend magnitude (km2):	
2.4.7	Population trend period:	1992-2003
2.4.8	Reasons for reported trend:	Improved knowledge/more accurate data
		Natural processes
	and/or specify:	
2.4.9	Justification of % thresholds for trends (optional):	
2.4.10	Main pressures:	943 - collapse of terrain, landslide
		965 - predation
2.4.11	Threats	965 - predation
2,5	Habitat for the species in the biogeographic region or marine region	
2.5.1	Habitats for the species:	Laderas, sotobosques, piederriscos y taliscas en sustratos basálticos con suelo pro
2.5.2	Area estimation (km2):	75
2.5.3	Date of estimation:	2007
2.5.4	Quality of the data:	Moderate e.g. based on partial data with some extrapolation
2.5.5	Trend of the habitat:	
2.5.6	Trend period:	1970-2006
2.5.7	Reasons for reported trend:	NaturalProcesses
	Other (specify):	
2.6	Future prospects for the species:	Poor prospects - species likely to struggle unless conditions change
2.7	Complementary information	
2.7.1	Favourable reference range (km2):	50
2.7.2	Favourable reference population:	
2.7.3	Suitable habitat for the species (km2):	50
2.7.4	Other relevant information (optional):	Favourable reference population: 150 individuals/population Poblaciones con fluc
Conclusion	Biogeographical or marine level	Conclusions within Natura 2000 sites (optional)
Conclusions: (2.3) Range:	Inadequate (U1)	
Conclusions: (2.4) Population:	Inadequate (U1)	
Conclusions: (2.5) Habitat for the species:	Unknown (XX)	

Sideritis discolor

Conclusions: (2.6) Future prospects:

Bad (U2)

Conclusions: Overall assessment:

Inadequate (U1)