



12 November 2015 Issue 435

Subscribe to free weekly News Alert

Source: Boersma, K.F., Vinken, G.C.M. & Jean Tournadre, J. (2015) Ships going slow in reducing their NOx emissions: changes in 2005-2012 ship exhaust inferred from satellite measurements over Europe. Environmental Research Letters, 10: 074007. DOI:10.1088/1748-9326/10/7/074007. This study is free to view at: http://iopscience.iop.org/ar ticle/10.1088/1748-9326/10/7/074007; jsessio nid=A289ABDCFC7CBC63A 561176A3762074C.c1

Contact:

folkert.boersma@wur.nl

Read more about: <u>Air pollution</u>, <u>Climate</u>

<u>change and energy,</u> <u>Environment and</u> <u>health, Sustainable</u> <u>business</u>

The contents and views included in Science for Environment Policy are based on independent, peer-reviewed research and do not necessarily reflect the position of the European Commission.

To cite this article/service: <u>"Science</u> for Environment Policy": European Commission DG Environment News Alert Service, edited by SCU, The University of the West of England, Bristol.

1. Smith, T.W.P. et al. (2014b) Third IMO GHG Study 2014. London: International Maritime Organisation.

Science for Environment Policy

Travelling slower reduces fuel consumption and nitrogen oxides emissions of ships

Ships that reduce their speed use less fuel, which lowers costs for shipping companies. The slow steaming practice also cuts nitrogen oxides (NO_x) emissions. A study found that ships travelling on four European routes lowered their NO_x emissions by 12% during the economic crisis of 2008/2009. Shipping continues to be a major way of transporting goods, however, and as the global economy recovers the researchers and civil society call for additional measures to reduce NO_x emissions from shipping and improve air quality in Europe.

Nitrogen oxides (NO_x = NO + NO₂) are toxic <u>air pollutants</u> and <u>greenhouse</u> gases that are released in large quantities from human activities, such as the combustion of fossil fuels. They also react with other compounds in the air to form further products, including particulate matter and smog, which harm <u>human health</u> and the environment.

Shipping contributes around 15% of global NO_x emissions¹. This study investigated how economic activity affects NO_x emissions, and in particular, the changes to ship operations introduced as a result of the 2008–2009 economic crisis.

The researchers estimated NO_x emissions from ships travelling along four European shipping lanes: the Mediterranean Sea, Bay of Biscay, North Sea, and Baltic Sea, for the period 2005–2012. They made their estimations using satellite observations of NO₂ from the Dutch-Finnish <u>Ozone Monitoring Instrument</u> (OMI) on-board <u>NASA's Aura satellite</u> combined with an atmospheric chemistry transport model. This type of model simulates the transport, chemistry, emissions and deposition of chemicals in the atmosphere.

They found that NO_x emissions increased by around 15% a year from 2005–2008, in line with increasing world trade, but then fell sharply by 12%, from 2008–2009. From 2009 to 2012, emissions remained steady at the 2009 level. These results mirror the impact of the economic crisis of 2008–2009 on shipping, which caused a dramatic reduction in the volume of goods being traded by sea.

Faced with falling revenues from lack of demand and overcapacity in the industry, <u>shipping</u> <u>companies</u> have sought ways to reduce their costs. One measure introduced from 2008 was to lower the speed at which ships travelled, to use less fuel. The practice, called slow steaming, reduces a ships speed from around 20–25 knots (approximately 37–46 km/hour) to 16–19 knots (~ 30–35 km/hour).

To confirm that ships were indeed going slower, the researchers calculated the average speed of ships travelling through the Mediterranean Sea from 2005–2012. They combined satellite readings of a ship's on-board Automatic Identification System, which tracks the vessel's location, with the number of ships entering or leaving the Suez Canal in Egypt, where the majority of ships pass, to calculate an average ship speed.

They found that, in the 2008–2009 period, ships travelled at a speed 30% slower than 2005 speeds. They also remained at this level of speed until 2012.

These results indicate that the shipping industry has widely adopted and continued to operate the voluntary practice of slow steaming since 2008/2009. Nevertheless, due to the steadily decreasing land emissions as a result of national and EU emission control policies, ship NO_x emissions as a share of total European NO_x emissions from all sources, including land-based emissions, has risen from 11% in 2005 to 14% in 2012.

Continued on next page.

Environment





12 November 2015 Issue 435

Subscribe to free weekly News Alert

Boersma, K.F., Vinken, G.C.M. & Jean Tournadre, J. (2015) Ships going slow in reducing their NO_x emissions: changes in 2005–2012 ship exhaust inferred from satellite measurements over Europe. Environmental Research Letters. 10: 074007. DOI:10.1088/1748-9326/10/7/074007. This study is free to view at: http://iopscience.iop.org/ar ticle/10.1088/1748-9326/10/7/074007;jsessio nid=A289ABDCFC7CBC63A

Contact: folkert.boersma@wur.nl

561176A3762074C.c1

Read more about:

<u>Air Pollution, Climate</u> <u>change and energy,</u> <u>Environment and</u> <u>health, Sustainable</u> business

The contents and views included in Science for Environment Policy are based on independent, peer-reviewed research and do not necessarily reflect the position of the European Commission.

To cite this article/service: <u>"Science</u> for Environment Policy": European Commission DG Environment News Alert Service, edited by SCU, The University of the West of England, Bristol.

Science for Environment Policy

Travelling slower reduces fuel consumption and nitrogen oxides emissions of ships (continued)

The researchers suggest that, in spite of the implementation of slow steaming, the shipping sector is responsible for an ever-larger share of European NO_x emissions and further policy measures are needed to realise reduction rates in ship emissions, similar to reductions seen in the road transport and energy generation sectors. It is likely that European shipping NO_x emissions will continue to grow as the global economy recovers, despite ships travelling slower, as shipping is a major way to transport goods.

Better monitoring of ship NO_x emissions, however, will soon be possible with the launch of new satellites with space-borne sensors, such as <u>Sentinel 4</u>, which will monitor air quality, stratospheric ozone and solar radiation, and be used in climate monitoring and to support European policies.





Environment