



MINISTERIO
DE MEDIO AMBIENTE, Y
MEDIO RURAL Y MARINO



ANEXO VII. BIBLIOGRAFÍA CONSULTADA.

PROLIFERACIONES

PROLIFERACIONES DE MEDUSAS

PELAGIA NOCTILUCA

MNEMIOPSIS LEIDYI

CARYBDEA MARSUPIALIS

RHIZOSTOMA PULMO

COTYLORHIZA TUBERCULATA



PROLIFERACIONES

Bay-Nouailhat A (2006) Description of *Thetys vagina* [on-line].

<http://www.european-marine-life.org/32/thetys-vagina.php>.

Boero F, Bouillon J, Gravili C, Miglietta MP, Parsons T & Piraino S (2008) Gelatinous plankton: irregularities rule the world (sometimes). *Marine Ecology Progress Series*, 356: 299-310, DOI: 10.3354/meps07368.

Boero F, Putti M, Trainito E, Prontera E, Piraino S & Shiganova TA (2009) First records of *Mnemiopsis leidyi* (Ctenophora) from the Ligurian, Thyrrenian and Ionian Seas (Western Mediterranean) and first record of *Phyllorhiza punctata* (Cnidaria) from the Western Mediterranean. *Aquatic Invasions*, 4 (4):675-680, DOI: 10.339/ai.2009.4.4.13.

Brodeur RD, Sugisaki H & Hunt GL (2002) Increases in jellyfish biomass in the Bering Sea: implications for the ecosystem. *Marine Ecology Progress Series*, 233: 89-103.

Gili, JM & Pagés, F (2005) Jellyfish blooms. *Bolletí de la Societat d'Història Natural de les Balears*, 48: 16-22.

León ME, Tejera E, Hernández F & Conway D (2005) Medusas de las islas de Cabo Verde: resultados de la campaña TFMCBM/98 Cabo Verde. *Vieraea*, 33: 11-28.

Molinero JC, Ibanez F, Nival P, Buecher E & Souissi S (2005) north Atlantic Climate and northwestern Mediterranean Plankton Variability. *Limnology and Oceanography*, 50 (4): 1213-1220.

Purcell JE (2005) Climate effects on formation of jellyfish and ctenophore blooms: a review. *Journal of the Marine Biological Association U.K.*, 85: 461-476.

Purcell JE, Hoover RA & Schwarck NT (2009) Interannual variation of strobilation by the scyphozoan *Aurelia labiata* in relation to polyp density, temperature, salinity and light conditions *in situ*. *Marine Ecology Progress Series*, 375: 139-149, DOI: 10.3354/meps07785.

Purcell JE, Uye S & Lo W (2007) Anthropogenic causes of jellyfish blooms and their direct consequences for humans: a review. *Marine Ecology Progress Series*, 350: 153-174, DOI: 10.3354/meps07093.

Richardson AJ, Bakun A, Hays GC & Gibbons MJ (2009) The jellyfish joyride: causes, consequences and management responses to a more gelatinous future. *Trends in Ecology and Evolution*, 24 (6): 312-322.



PROLIFERACIONES DE MEDUSAS

- Alvarez G, Benovic A, Malej A, Lucic D, Makovec T, Onofri V, Acha M, Madirolas A & Mianzan H (2009) Acoustic survey of a jellyfish-dominated ecosystem (Mljet Island, Croatia). *Hydrobiologia*, 616: 99-111, DOI: 10.1007/s10750-008-9587-6.
- Arai MN (2001) Pelagic coelenterates and eutrophication: a review. *Hydrobiologia*, 451: 69-87.
- Arai MN (2009) The potential importance of podocysts to the formation of scyphozoan blooms: a review. *Hydrobiologia*, 616: 241-246, DOI: 10.1007/s10750-008-9588-5.
- Bajcar T, Malacic V, Malej A & Sirok B (2009) Kinematic properties of the jellyfish *Aurelia* sp. *Hydrobiologia*, 616: 279-289, DOI: 10.1007/s10750-008-9589-4.
- Baxter JE, Walne AW, Purcell JE, McAllen R & Doyle TK (2010) Identification of jellyfish from Continuous Plankton Recorder samples. *Hydrobiologia*, DOI: 10.1007/s10750-010-0212-0.
- Bayha KM & Graham WM (2009) A new Taqman® PCR-based method for the detection and identification of scyphozoan jellyfish polyps. *Hydrobiologia*, 616: 217-228, DOI: 10.1007/s10750-008-9590-y.
- Burnett JW (2001) Medical aspects of jellyfish envenomation: pathogenesis, case reporting and therapy. *Hydrobiologia*, 451: 1-9.
- Condon RH, Decker MB & Purcell JE (2001) Effects of low dissolved oxygen on survival and asexual reproduction of scyphozoan polyps (*Chrysaora quinquecirrha*). *Hydrobiologia*, 451: 89-95.
- D'Ambra I, Costello JH & Bentivegna F (2001) Flow and prey capture by the scyphomedusa *Phyllorhiza punctata* von Lendenfeld, 1884. *Hydrobiologia*, 451: 223-227.
- Dawson MN & Martin KE (2001) Geographic variation and ecological adaptation in *Aurelia* (Scyphozoa, Semaestomaeae): some implications from molecular phylogenetics. *Hydrobiologia*, 451: 229-273.
- Dawson MN, Martin LE & Penland LK (2001) Jellyfish swarms, tourists, and the Christ-child. *Hydrobiologia*, 451: 131-144.
- Dong J, Jiang L, Tan K, Liu H, Purcell JE, Li P & Ye C (2009) Stock enhancement of the edible jellyfish (*Rhopilema esculentum* Kishinouye) in Liaodong Bay, China: a review. *Hydrobiologia*, 616: 113-118, DOI: 10.1007/s10750-008-9592-9.



- Finenko GA, Anninsky BE, Romanova ZA, Abolmasova GI & Kideys AE (2001) Chemical composition, respiration and feeding rates of the new alien ctenophore, *Beroe ovata*, in the Black Sea. *Hydrobiologia*, 451: 177-186.
- Friis L, Martínez J & Tiselius P (2010) Bioenergetics and growth in the ctenophore *Pleurobrachia pileus*. *Hydrobiologia*, DOI: 10.1007/s10750-010-0219-6.
- Frost JR, Jacoby CA & Youngbluth MJ (2010) Behaviour of *Nemopsis bachei* L. Agassiz, 1849 medusae in the presence of physical gradients and biological thin layers. *Hydrobiologia*, 645: 97-111, DOI: 10.1007/s10750-010-0213-z.
- Fuentes V, Angel DL, Bayha KM, Atienza D, Edelist D, Bordehore C, Gili JM & Purcell JE (2010) Blooms of the invasive ctenophore, *Mnemiopsis leidyi*, span the Mediterranean Sea in 2009. *Hydrobiologia*, DOI: 10.1007/s10750-010-0205-z.
- Gibbons MJ & Richardson AJ (2009) Patterns of jellyfish abundance in the north Atlantic. *Hydrobiologia*, 616: 51-65, DOI: 10.1007/s10750-008-9593-8.
- Gordon MR & Seymour JE (2009) Quantifying movement of the tropical Australian cubozoan *Chironex fleckeri* using acoustic telemetry. *Hydrobiologia*, 616: 87-97, DOI: 10.1007/s10750-008-9594-7.
- Graham WM (2001) Numerical increases and distributional shifts of *Chrysaora quinquecirrha* (Desor) and *Aurelia aurita* (Linné) (Cnidaria: Scyphozoa) in the northern Gulf of Mexico. *Hydrobiologia*, 451: 97-111.
- Graham WM, Pagès F & Hammer WM (2001) A physical context for gelatinous zooplankton aggregation: a review. *Hydrobiologia*, 451: 199-212.
- Hammer WM & Dawson MN (2009) A review and synthesis on the systematics and evolution of jellyfish blooms: advantageous aggregations and adaptive assemblages. *Hydrobiologia*, 616: 161-191, DOI: 10.1007/s10750-008-9620-9.
- Helmholz H, Johnston BD, Ruhnau C & Prange A (2010) Gill cell toxicity of northern boreal scyphomedusae *Cyanea capillata* and *Aurelia aurita* measured by an in vitro cell assay. *Hydrobiologia*, DOI: 10.1007/s10750-010-0216-9.
- Holst S & Jarms G (2010) Effects of low salinity on settlement and strobilation of scyphozoa (Cnidaria): Is the lion's mane *Cyanea capillata* (L.) able to reproduce in the brackish Baltic Sea? *Hydrobiologia*, DOI: 10.1007/s10750-010-0214-y.
- Hoover R & Purcell JE (2009) Substrate preferences of scyphozoan *Aurelia labiata* polyps among common dock-building materials. *Hydrobiologia*, 616: 259-267, DOI: 10.1007/s10750-008-9595-6.



- Hsieh YP, Leong F & Rudloe J (2001) Jellyfish as food. *Hydrobiologia*, 451:11-17.
- Ishii H & Tanaka F (2001) Food and feeding of *Aurelia aurita* in Tokyo Bay with an analysis of stomach contents and a measurement of digestion times. *Hydrobiologia*, 451: 311-320.
- Johnson DR, Perry HM & Burke D (2001) Developing jellyfish hypotheses using circulation models. *Hydrobiologia*, 451: 213-221.
- Ki J, Kim I & Lee J (2009) Comparative analysis of nuclear ribosomal DNA from the moon jelly *Aurelia* sp.1 (Cnidaria: Scyphozoa) with characterizations of the 18S, 28S genes, and the intergenic spacer (IGS). *Hydrobiologia*, 616: 229-239, DOI: 10.1007/s10750-008-9596-5.
- Kogovsek T, Bogunovic B & Malej A (2010) Recurrence of bloom-forming scyphomedusae: wavelet analysis of a 200-year time series. *Hydrobiologia*, DOI: 10.1007/s10750-010-0217-8.
- Lassen S, Helmholz H, Ruhnau C & Prange A (2010) Characterisation of neurotoxic polypeptides from *Cyanea capillata* medusae (Scyphozoa). *Hydrobiologia*, 645: 213-221, DOI: 10.1007/s10750-010-0221-z.
- Liu W, Lo W, Purcell JE & Chang H (2009) Effects of temperature and light intensity on asexual reproduction of the scyphozoan, *Aurelia aurita* (L.) in Taiwan. *Hydrobiologia*, 616: 247-258, DOI: 10.1007/s10750-008-9597-4.
- Lucas CH (2001) Reproduction and life history strategies of the common jellyfish, *Aurelia aurita*, in relation to its ambient environment. *Hydrobiologia*, 451: 229-246.
- Martorelli SR (2001) Digenea parasites of jellyfish and ctenophores of the Southern Atlantic. *Hydrobiologia*, 451: 305-310.
- Masuda R (2009) Ontogenetic changes in the ecological function of the association behaviour between jack mackerel *Trachurus japonicus* and jellyfish. *Hydrobiologia*, 616: 269-277, DOI: 10.1007/s10750-008-9598-3.
- Mianzan H, Martos P, Costello JH & Guerrero RA (2010) Avoidance of hydrodynamically mixed environments by *Mnemiopsis leidyi* (Ctenophora: Lobata) in open-sea populations from Patagonia, Argentina. *Hydrobiologia*, DOI: 10.1007/s10750-010-0218-7.
- Mills CE (2001) Jellyfish blooms: are populations increasing globally in response to changing ocean conditions? *Hydrobiologia*, 451: 55-68.
- Moss AG, Estes AM, Muellner LA & Morgan DD (2001) Protistan epibionts of the ctenophore *Mnemiopsis mccradyi* Mayer. *Hydrobiologia*, 451: 295-304.



- Moss AG, Rapoza RC & Muellner L (2001) A novel cilia-based feature within the food grooves of the ctenophore *Mnemiopsis mccradyi* Mayer. *Hydrobiologia*, 451: 287-294.
- Omori M & Nakano E (2001) Jellyfish fisheries in Southeast Asia. *Hydrobiologia*, 451: 19-26.
- Pauly D, Graham W, Libralato S, Morissette L & Palomares MLD (2009) Jellyfish in ecosystems, online databases, and ecosystem models. *Hydrobiologia*, 616: 67-85, DOI: 10.1007/s10750-008-9583-x.
- Pitt KA, Connolly RM & Meziane T (2009) Stable isotope and fatty acid tracers in energy and nutrient studies of jellyfish: a review. *Hydrobiologia*, 616: 119-132, DOI: 10.1007/s10750-008-9581-z.
- Pitt KA, Welsh DT & Condon RH (2009) Influence of jellyfish blooms on carbon, nitrogen and phosphorus cycling and plankton production. *Hydrobiologia*, 616: 133-149, DOI: 10.1007/s10750-008-9584-9.
- Purcell JE & Arai MN (2001) Interactions of pelagic cnidarians and ctenophores with fish: a review. *Hydrobiologia*, 451: 27-44.
- Purcell JE (2009) Extension of methods for jellyfish and ctenophore trophic ecology to large-scale research. *Hydrobiologia*, 616: 23-50, DOI: 10.1007/s10750-008-9585-8.
- Purcell JE, Fuentes V, Atienza D, Tilves U, Astorga D, Kawahara M & Hays GC (2010) Use of respiration rates of scyphozoan jellyfish to estimate their effects on the food web. *Hydrobiologia*, DOI: 10.1007/s10750-010-0240-9.
- Purcell JE, Shiganova TA, Decker MB & Houde ED (2001) The ctenophore *Mnemiopsis* in native and exotic habitats: U.S. estuaries *versus* the Black Sea basin. *Hydrobiologia*, 451: 145-176.
- Quiñones J, González V, Zeballos J, Purca S & Mianzan H (2010) Effects of El Niño-driven environmental variability on black turtle migration to Peruvian foraging grounds. *Hydrobiologia*, DOI: 10.1007/s10750-010-0225-8.
- Raskoff KA (2001) The impact of El Niño events on populations of mesopelagic hydromedusae. *Hydrobiologia*, 451: 121-129.
- Sabatés A, Pagès F, Atienza D, Fuentes V, Purcell JE & Gili JM (2010) Planktonic cnidarian distribution and feeding of *Pelagia noctiluca* in the NW Mediterranean Sea. *Hydrobiologia*, 645: 153-165, DOI: 10.1007/s10750-010-0221-z.



- Sexton MA, Hood RR, Sarkodee-adoo J & Liss AM (2010) Response of *Chrysaora quinquecirrha* medusae to low temperature. *Hydrobiologia*, DOI: 10.1007/s10750-010-0222-y.
- Shiganova TA, Bulgakova YV, Volovik SP, Mirzoyan ZA & Dudkin SI (2001) The new invader *Beroe ovata* Mayer 1912 and its effect on the ecosystem in the northeastern Black Sea. *Hydrobiologia*, 451: 187-197.
- Sparks C, Buecher E, Brierley AS, Axelsen BE, Boyer H & Gibbons MJ (2001) Observations on the distribution and relative abundance of the scyphomedusan *Chrysaora hysoscella* (Linné, 1766) and the hydrozoan *Aequorea aequorea* (Forskål, 1755) in the northern Benguela ecosystem. *Hydrobiologia*, 451: 275-286.
- Straehler-Pohl I & Jarms G (2010) Identification key for young ephyrae: a first step for early detection of jellyfish blooms. *Hydrobiologia*, 645: 3-21, DOI: 10.1007/s10750-010-0226-7.
- Sullivan BK, Van Keuren D & Clancy M (2001) Timing and size of blooms of the ctenophore *Mnemiopsis leidyi* in relation to temperature in Narragansett Bay, RI. *Hydrobiologia*, 451: 113-120.
- Tinta T, Malej A, Kos M & Turk V (2010) Degradation of the Adriatic medusa *Aurelia* sp. by ambient bacteria. *Hydrobiologia*, 645: 179-191, DOI: 10.1007/s10750-010-0223-x.
- Uye S (2010) Human forcing of the copepod-fish-jellyfish triangular trophic relationship. *Hydrobiologia*, DOI: 10.1007/s10750-010-0208-9.
- Watanabe T & Ishii H (2001) *In situ* estimation of ephyrae liberated from polyps of *Aurelia aurita* using settling plates in Tokyo Bay, Japan. *Hydrobiologia*, 451: 247-258.
- West EJ, Welsh DT & Pitt KA (2009) Influence of decomposing jellyfish on the sediment oxygen demand and nutrient dynamics. *Hydrobiologia*, 616: 151-160, DOI: 10.1007/s10750-008-9586-7.
- Wiebring A, Helmholz H, Lassen S, Prange A & Jarms G (2010) Separation and analysis of different types of nematocysts from *Cyanea capillata* (L.) medusae. *Hydrobiologia*, DOI: 10.1007/s10750-010-0227-6.
- Winans AK & Purcell JE (2010) Effects of pH on asexual reproduction and statolith formation of the scyphozoan, *Aurelia labiata*. *Hydrobiologia*, DOI: 10.1007/s10750-010-0224-9.





MINISTERIO
DE MEDIO AMBIENTE, Y
MEDIO RURAL Y MARINO

Youngbluth MJ & Båmstedt U



(2001) Distribution, abundance, behaviour and metabolism of *Periphylla periphylla*, a mesopelagic coronate medusa in a Norwegian fjord. *Hydrobiologia*, 451: 321-333.

PELAGIA NOCTILUCA

- Goy J, Morand P & Etienne M (1989) Long-term fluctuations of *Pelagia noctiluca* (Cnidaria, Scyphomedusa) in the western Mediterranean Sea. Prediction by climatic variables. *Deep-Sea Research*, 36 (2): 269-279.
- Licandro P, Conway DVP, Daly MN, Fernández ML, Gasparini S, Hecq JH, Tranter P & Kirby RR (2010) A blooming jellyfish in the northeast Atlantic and Mediterranean. *Biology Letters*, DOI: 10.1098/rsbl.2010.0150.
- Mariottini GL & Pane L (2010) Mediterranean Jellyfish Venoms: A Review on Scyphomedusae. *Marine Drugs*, 8: 1122-1152, DOI: 10.3390/md8041122.
- Mariottini GL, Giacco E & Pane L (2008) The Mauve Stinger *Pelagia noctiluca* (Forsskål, 1775). Distribution, Ecology, Toxicity and Epidemiology of Stings. A Review. *Marine Drugs*, 6: 496-513, DOI: 10.3390/md20080025.
- Rottini L & Avian M (1991) Reproduction of *Pelagia noctiluca* in the central and northern Adriatic Sea. *Hydrobiologia*, 216/217: 197-202.
- Sabatés A, Pagès F, Atienza D, Fuentes V, Purcell JE & Gili JM (2010) Planktonic cnidarian distribution and feeding of *Pelagia noctiluca* in the NW Mediterranean Sea. *Hydrobiologia*, 645: 153-165, DOI: 10.1007/s10750-010-0221-z.

MNEMIOPSIS LEIDYI

- Fuentes V, Angel DL, Bayha KM, Atienza D, Edelist D, Bordehore C, Gili JM & Purcell JE (2010) Blooms of the invasive ctenophore, *Mnemiopsis leidyi*, span the Mediterranean Sea in 2009. *Hydrobiologia*, DOI: 10.1007/s10750-010-0205-z.
- Fuentes V, Atienza D, Gili JM & Purcell JE (2009) First records of *Mnemiopsis leidyi* A. Agassiz 1865 off the NW Mediterranean coast of Spain. *Aquatic Invasions*, 4 (4): 671-674, DOI: 10.3391/ai.2009.4.4.12.
- Purcell JE, Shiganova TA, Decker MB & Houde ED (2001) The ctenophore *Mnemiopsis* in native and exotic habitats: U.S. estuaries versus the Black Sea basin. *Hydrobiologia*, 451: 145-176.



Ulrik H, Bøttiger L, Madsen CV & Purcell JE (2007) Invasive ctenophore *Mnemiopsis leidyi* in Limfjorden (Denmark) in late summer 2007-assessment of abundance and predation effects. *Aquatic Invasions*, 2 (4): 395-401, DOI: 10.3391/ai.2007.2.4.8.

CARYBDEA MARSUPIALIS

De Donno A, Idolo A & Bagordo F (2009) Epidemiology of jellyfish stings reported to summer health centres in the Salento peninsula (Italy). *Contact Dermatitis*, 60: 330-335.

Di Camillo D, Bo M, Puce S, Tazioli S & Bavestrello G (2006) The cnidome of *Carybdea marsupialis* (Cnidaria: Cubomedusae) from the Adriatic Sea. *Journal of the Marine Biological Association*, 86: 705-709.

Kazmi QB & Sultana R (2008) *Carybdea marsupialis* (Cnidaria: Cubomedusae)-observed for the first time in Gwadar Bay Waters (Arabian Sea). *Marine Biodiversity Records*, 1: e80, DOI: 10.1017/S1755267207008251.

Straehler-Pohl I & Jarms G (2005) Life cycle of *Carybdea marsupialis* Linnaeus, 1758 (Cubozoa, Carybdeidae) reveals metamorphosis to be a modified strobilation. *Marine Biology*, DOI: 10.1007/s00227-005-0031-4.

PHYSALIA PHYSALIS

Burnett JW, Ordonez JV & Calton GJ (1986) Differential toxicity of *Physalia physalis* (Portuguese man-o'-war) nematocysts separated by flow cytometry. *Toxicon*, 24 (5): 514-518.

Calton GJ & Burnett JW (1973) The purification of Portuguese man-of-war nematocyst toxins by gel diffusion. *Comparative and General Pharmacology*, 4: 267-270.

Copeland DE (1968) Fine structures of the carbon monoxide secreting tissue in the float of Portuguese man-of-war *Physalia physalis* L. *Biological Bulletin*, 135 (3): 486-500.

Dunn CW & Wagner GP (2006) The evolution of colony-level development in the Siphonophora (Cnidaria: Hydrozoa). *Development Genes and Evolution*, 216:743-754.



- Fagetti E (1958) Nota sobre *Physalia physalis* L. procedente de la Isla de Pascua. *Revista de Biología Marina*, 7 (1,2 y 3): 188-190.
- Fenner P (1997) Awareness, Prevention and Treatment of world-wide marine stings and bites. International Life Saving Federation Medical/Rescue Conference Proceedings [on-line].
http://www.usla.org/PublicInfo/library/Treatment_of_Marine_Stings.pdf
- Iosilevskii G & Weish D (2009) Hydrodynamics of sailing of the Portuguese man-of-war *Physalia physalis*. *Journal of the Royal Society Interface*, 6: 613-626, DOI: 10.1098/RSIF.2008.0457.
- Lane CE & Dodge E (1958) The toxicity of *Physalia* nematocysts. *The Biological Bulletin*, 115: 219-226.
- Martínez M, Villena ME, Marín I & Monedero J (2010) Un paciente con... picadura por Carabela Portuguesa, una "medusa" algo especial. *Revista Clínica de Medicina de Familia*, 3 (2): 143-145.
- Palma S (1999) Sifonóforos (Cnidaria, Hydrozoa) de aguas superficiales de Isla de Pascua. *Investigaciones Marinas*, 27: 19-23.
- Purcell JE (1984) Predation on fish larvae by *Physalia physalis*, the Portuguese man of war. *Marine Ecology Progress Series*, 19: 189-191.
- Southeastern Regional Taxonomic Center. South Carolina Department of Natural Resources (2010). The Portuguese man-of-war [on-line].
<http://www.dnr.sc.gov/marine/sertc/>
- Tamkun MM & Hessinger DA (1981) Isolation and partial characterization of a hemolytic and toxic protein from the nematocyst venom of the Portuguese Man-of-war, *Physalia physalis*. *Biochimica et Biophysica Acta*, 667: 87-98.
- Wittenberg JB, Noronha JM & Silverman M (1962) Folic Acid Derivates in the Gas Gland of *Physalia physalis* L. *Biochemistry Journal*, 85: 9-15.

RHIZOSTOMA PULMO

- Kokelj F & Plozzer C (2002) Irritant contact dermatitis from the jellyfish *Rhizostoma pulmo*. *Contact Dermatitis*, 46 (3): 179-180.
- Mariottini GL & Pane L (2010) Mediterranean Jellyfish Venoms: A Review on Scyphomedusae. *Marine Drugs*, 8: 1122-1152, DOI: 10.3390/md8041122.





MINISTERIO
DE MEDIO AMBIENTE, Y
MEDIO RURAL Y MARINO



Muhammed F & Sultana R

(2008) New record of edible jellyfish, *Rhizostoma pulmo* (Cnidaria: Scyphozoa: Rhizostomidae) from Pakistani waters. Marine Biodiversity Records, 1e: 67, DOI: 10.1017/S1755267207007294.

COTYLORHIZA TUBERCULATA

Kikinger R (1992) *Cotylorhiza tuberculata* (Cnidaria: Scyphozoa)- Life History of a Stationary Population. Marine Ecology, 13 (4): 333-362.

Stopar K, Ramsak A, Trontelj P & Malej A (2010) Lack of genetic structure in the jellyfish *Pelagia noctiluca* (Cnidaria: Scyphozoa: Semaestomeae) across European seas. Molecular Phylogenetics and Evolution, 57 (1): 417-428.

